



## Lecture 7 : Second-order methods - Newton algorithm

This lecture is dedicated to the second-order methods, notably the Newton algorithms and its theoretical guarantee. In general, these methods can enjoy a much faster convergence rate than first-order counterparts. Nevertheless, the cost per iteration makes them very expensive, hence hindering their usage in large-scale optimization. Adapting second-order methods to real-world scale problems is an active research domain, and might enable us to handle challenging optimization problems.

### 1 Second-order methods - the Newton step

So far, all algorithms that we have seen in this course only exploit first-order information, i.e., their update rules only use gradients. It is, thus, very tempting to exploit higher-order derivatives to see whether we can break the fundamental limit of first-order methods. Newton method, possibly one of the most well-known second-order methods, will be our focus of today lecture.

Given a  $C^2$  function, the Newton update is given by:

$$x_{k+1} = x_k - \alpha \nabla^2 f(x_k)^{-1} \nabla f(x_k). \quad (\text{Newton})$$

When  $\alpha = 1$ , it is called the *pure Newton step*. If the step-size  $\alpha \neq 1$ , then this is a damped Newton step.

There are, in fact, many interpretations and fun facts about (Newton):

**Original Newton method is to find zero of functions, and not an optimization algorithm** The original Newton method was proposed to find the  $x$  such that  $f(x) = 0$  of a differentiable function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Starting from a guess point, the algorithm updates its iterations as:

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}.$$

This rule is proposed due to the first-order approximation:  $f(x + \delta) = f(x) + f'(x)\delta$ . The pure (Newton) is when we apply the original Newton method for  $f : \mathbb{R}^n \rightarrow \mathbb{R}^n : x \mapsto \nabla f(x)$ , or more precisely:

$$\nabla^2 f(x_k)(x_{k+1} - x_k) - \nabla f(x_k) = 0.$$

**Pure Newton step is the steepest descent method of a special metric** In fact, the descent direction of gradient descent can be view as the minimizer of the following constrained problem:

$$\begin{aligned} \underset{p \in \mathbb{R}^d}{\text{Minimize}} \quad & \nabla f(x_k)^\top p \\ \text{s.t.} \quad & \|p\|_2 \leq 1. \end{aligned}$$

The Newton step has the same interpretation, but with  $\|p\|_f = p^\top \nabla^2 f(x_k) p$ , i.e.,

$$\begin{aligned} \underset{p \in \mathbb{R}^d}{\text{Minimize}} \quad & \nabla f(x_k)^\top p \\ \text{s.t.} \quad & \|p\|_f \leq 1. \end{aligned}$$

We can get the optimal direction by considering  $\nabla^2 f^{1/2} p$ .

**Pure Newton step is the minimization of the surrogate of the function  $f$**  In fact, the gradient descent step can be viewed as the minimizer of the following surrogate function:

$$f_{\text{GD}}(x) = f(x_k) + \nabla f(x_k)^\top (x - x_k) + \frac{1}{2\alpha} \|x - x_k\|^2$$

where the minimizer is given by:

$$x^* = x_k - \alpha \nabla f(x_k) = \text{gradient descent step.}$$

For Newton step, the surrogate function is the second order Taylor approximation of  $f$ , i.e.:

$$f_{\text{Newton}}(x) = f(x_k) + \nabla f(x_k)^\top (x - x_k) + \frac{1}{2} (x - x_k)^\top \nabla^2 f(x_k) (x - x_k),$$

given that  $\nabla^2 f(x_k) \succ 0$ .

**Affine invariant of Newton step** An important of (Newton) is that it is independent of linear or affine changes of coordinates. Suppose that we have a non-singular matrix  $\mathbf{A} \in \mathbb{R}^{n \times n}$ , and define  $g = f \circ \mathbf{A}$ , we have:

$$\nabla g(x) = \mathbf{A}^\top \nabla f(\mathbf{A}x) \quad , \quad \nabla^2 g(x) = \mathbf{A}^\top \nabla^2 f(\mathbf{A}x) \mathbf{A}.$$

Therefore, the Newton update of the function  $g$  at the point  $y = \mathbf{A}^{-1}x$  is:

$$\begin{aligned} \mathbf{A}^{-1}x - (\nabla^2 g(\mathbf{A}^{-1}x))^{-1} \nabla g(\mathbf{A}^{-1}x) &= x - (\mathbf{A}^\top \nabla^2 f(x) \mathbf{A})^{-1} \mathbf{A}^\top \nabla f(x) \\ &= \mathbf{A}^{-1}x - \mathbf{A}^{-1} \nabla^2 f(x)^{-1} \nabla f(x) \\ &= \mathbf{A}^{-1}(x - \nabla^2 f(x)^{-1} \nabla f(x)). \end{aligned}$$

Thus, the Newton update of the function  $g$  is that of the function  $f$ , up to the inverse of non-singular linear transformation  $\mathbf{A}$ . This is the unique property of Newton methods, unlike other first-order methods.

## 2 Newton algorithm and theoretical guarantees

The Newton algorithm is given as below: in each iteration:

1. Compute the Newton step  $p_k$  as in (Newton) and decrement  $\lambda^2 := \nabla f(x)^\top \nabla^2 f(x)^{-1} \nabla f(x)$ .
2. Line search: find step-size  $\alpha$  so that  $f(x - \alpha p_k) \leq f(x) - c\alpha \lambda^2$  (Armijo rule) by backtracking.
3. Update:  $x_{k+1} = x_k - \alpha p_k$ .

**Theoretical guarantees** We employ two assumptions

**Assumption 2.1** (Strongly convex and smooth). *The function  $f$  is  $m$ -strongly convex and  $M$ -smooth, i.e.,*

$$m\mathbf{I} \leq \nabla^2 f(x) \leq M\mathbf{I}, \forall x.$$

**Assumption 2.2** ( $L$ -Lipschitz hessian). *The Hessian of  $f$  is Lipschitz continuous with constant  $L$ , i.e.,*

$$\|\nabla^2 f(x) - \nabla^2 f(y)\|_2 \leq L\|x - y\|_2, \forall x, y.$$

In the analysis, the Newton algorithm will go through two stage:

1. *Damped Newton phase*: In this stage, the value function will decrease steadily by a constant  $\gamma > 0$  and the norm of  $\|\nabla f(x)\| \leq \delta > 0$ .
2. *Pure Newton phase*: In the second stage, the norm of the gradient becomes small and the line search will always pick  $\alpha_k = 1$  and:

$$\frac{L}{2m} \|\nabla f(x_{k+1})\|_2 \leq \left( \frac{L}{2m} \|\nabla f(x_k)\|_2 \right)^2.$$

This is also known as quadratic convergence.

Before entering the technicalities, remind us of two identities that will be used repeatedly throughout the proof:

$$\begin{aligned} f(y) &= f(x) + \int_0^1 \nabla f(x + t(y-x))(y-x) dt, \\ f(y) &= f(x) + \nabla f(x)^\top (y-x) + \int_0^1 (1-t)(y-x)^\top \nabla^2 f(x + t(y-x))(y-x) dt. \end{aligned} \tag{1}$$

**Damped Newton phase** By  $M$ -smoothness of the function  $f$ , we have:

$$\begin{aligned} f(x_k - \alpha p_k) &\leq f(x_k) - \alpha \nabla f(x_k)^\top p_k + \alpha^2 \frac{M \|p_k\|^2}{2} \\ &\leq f(x_k) - \alpha \lambda_k^2 + \frac{M}{2m} \alpha^2 \lambda_k^2. \end{aligned}$$

Therefore, if we choose  $\alpha \leq \frac{m}{M}$ , it will satisfy the exit condition of the line search for  $c < \frac{1}{2}$  because:

$$f(x_k - \alpha p_k) \leq f(x_k) - \alpha \left( 1 - \frac{M}{2m} \alpha \right) \lambda^2. \tag{2}$$

Therefore, our step-size  $\alpha$  found by line-search will satisfy:

$$\alpha_k \geq \beta \frac{m}{M},$$

where  $\beta > 0$  is a constant dependent on our line search algorithm. As such:

$$f(x_{k+1}) - f(x) \leq -\frac{1}{2} \alpha \lambda^2 \leq -\beta \frac{m}{2M} \lambda^2 \leq -\alpha \beta \eta^2 \frac{m}{M^2}.$$

because  $\|\nabla f(x)\|_2 < \eta$  and  $\lambda^2 \geq (1/M) \|\nabla f(x)\|^2$ .

**Quadratically convergent phase** We prove next that if:

$$\eta \leq \frac{m^2}{2L},$$

then the line search will choose unit step, i.e.,  $\alpha_k = 1$ . Indeed, since  $f$  is  $L$ -smooth, we have:

$$\begin{aligned} f(x_{k+1}) &= f(x_k - \underbrace{\nabla^2 f(x_k)^{-1} \nabla f(x_k)}_{p_k}) \\ &= f(x_k) - \nabla f(x_k)^\top p_k + \int_0^1 (1-t) p_k^\top \nabla^2 f(x_k - tp_k) p_k dt \\ &= f(x_k) - \frac{1}{2} \lambda_k^2 + \int_0^1 (1-t) p_k^\top (\nabla^2 f(x_k - tp_k) - \nabla^2 f(x_k)) p_k dt \\ &\leq f(x_k) - \frac{1}{2} \lambda_k^2 + \left\| \int_0^1 (1-t) p_k^\top (\nabla^2 f(x_k - tp_k) - \nabla^2 f(x_k)) p_k dt \right\|^2 \\ &\leq f(x_k) - \frac{1}{2} \lambda_k^2 + \frac{L}{6} \cdot \|p_k\|^3. \end{aligned}$$

Because

$$\begin{aligned} \lambda_k^2 &= \|\nabla^2 f(x_k)^{-\frac{1}{2}} \nabla f(x_k)\|_2^2 = \|\nabla^2 f(x_k)^{\frac{1}{2}} p_k\|_2^2 \geq m \|p_k\|^2, \\ \|\nabla f(x_k)\| &= \|\nabla^2 f(x_k) p_k\|_2 \geq m \|p_k\|, \end{aligned}$$

we have:

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2} \lambda_k^2 + \frac{L}{6m^2} \lambda_k^2 \|\nabla f(x_k)\| = f(x_k) - \frac{1}{2} \lambda_k^2 \left(1 - \frac{L \|\nabla f(x_k)\|}{3m^2}\right).$$

Therefore, if  $\eta \leq \frac{m^2}{2L}$ , we have  $f(x_k - d_k) \leq f(x_k) - \frac{1}{4} \lambda_k^2$  and line search will choose  $\alpha_k = 1$ .

Next, we will argue that when  $\|\nabla f(x_k)\| \leq \frac{m^2}{2L}$ ,  $\|\nabla f(x_\ell)\|$ ,  $\ell \geq k$  will be smaller than this amount forever. Indeed,

$$\begin{aligned} \|\nabla f(x_{k+1})\|_2 &= \|\nabla f(x_k - \nabla^2 f(x_k)^{-1} \nabla f(x_k))\|_2 \\ &= \left\| \nabla f(x_k) - \int_0^1 \nabla^2 f(x_k - t \nabla^2 f(x_k)^{-1} \nabla f(x_k)) \nabla^2 f(x_k)^{-1} \nabla f(x_k) dt \right\|_2 \\ &= \left\| \int_0^1 (\nabla^2 f(x_k - t \nabla^2 f(x_k)^{-1} \nabla f(x_k)) - \nabla^2 f(x_k)) \nabla^2 f(x_k)^{-1} \nabla f(x_k) dt \right\|_2 \\ &\leq \int_0^1 \|(\nabla^2 f(x_k - t \nabla^2 f(x_k)^{-1} \nabla f(x_k)) - \nabla^2 f(x_k)) \nabla^2 f(x_k)^{-1} \nabla f(x_k)\|_2 dt \\ &\leq L \int_0^1 t \|\nabla^2 f(x_k)^{-1} \nabla f(x_k)\| \|\nabla^2 f(x_k)^{-1}\| \|\nabla f(x_k)\|_2 dt \\ &\leq \frac{L}{2m^2} \|\nabla f(x_k)\|_2^2 \leq \|\nabla f(x_k)\|_2. \end{aligned}$$

Moreover, the above inequality also shows that:

$$\|\nabla f(x_k)\|_2 = O(c^{2^k}), \quad 0 \leq c < 1,$$

which is known as quadratic convergence speed of the norm of the gradient. This is much faster than linear convergence rate, which is read as:  $\|\nabla f(x_k)\|_2 = O(c^k)$ ,  $c \in [0, 1)$ . Finally, to get the quadratic convergence for the distance  $\|x_k - x^*\|$ , we have:

$$m \|x_k - x^*\|_2^2 \leq \langle \nabla f(x_k) - \nabla f(x^*), x_k - x^* \rangle \leq \|\nabla f(x_k)\| \|x_k - x^*\|.$$

Hence,

$$\|\nabla f(x_k)\| \geq m\|x_k - x^*\|_2.$$

Note that when one enters the quadratically convergent phase, to reach an error  $\epsilon$ , one needs roughly  $\log_2 \log_2 \frac{1}{\epsilon}$ . For machine precision error (approximately  $10^{-16}$ ), one practically needs at most 6 iterations. Therefore, the number of iterations of Newton method is upper-bounded by:

$$\frac{f(x_0) - f^*}{\eta} + 6.$$

The Newton method is suitable for small to medium sized problems since it suffers from a couple of problems:

1. Inverting the Hessian is expensive for large-scaled optimization problems since it might cost up to  $O(d^3)$  basic operations where  $d$  is the problem dimension. Without special structure of the Hessian, this complexity per iteration is prohibitive.
2. If the function  $f$  is not strongly convex, the Hessian is not invertible and renders the Newton method useless. There are a couple of workarounds such as:
  - (a) We choose the direction  $p_k$  as:

$$p_k \in \operatorname{argmin}_{p \in \mathbb{R}^d} \|\nabla^2 f(x_k)p - \nabla f(x_k)\|_2.$$

In case  $\nabla^2 f(x_k)$  is invertible, then we recover the Newton method. Otherwise, one needs to use a least square solver to find  $d_k$ .

- (b) We add a perturbation to the Hessian matrix and compute the direction as:

$$p_k = (\nabla^2 f(x_k) + \epsilon \mathbf{I})^{-1} \nabla f(x_k).$$

where  $\epsilon > 0$  is a hyper-parameter of the algorithm.

3. If the function is not (strongly) convex, then the Newton direction is not necessarily a descent direction, i.e.,

$$\lambda^2(x_k) := \nabla f(x_k)^\top \nabla^2 f(x_k)^{-1} \nabla f(x_k) \geq 0,$$

if  $\nabla f(x_k)$  is an eigenvector of  $\nabla^2 f(x_k)$  with a negative eigenvalue. Using a similar model to the Newton method, many authors replaced the exact hessian matrix by a semi-definite approximation  $\mathbf{B}_k$ . These methods are known as quasi-Newton methods and they update the iteration along the direction  $d$  defined as:

$$p_k \in \operatorname{argmin}_{p \in \mathbb{R}^d} f(x_k) + \nabla f(x_k)^\top p + \frac{1}{2} p^\top \mathbf{B}_k p.$$

If one chooses  $\mathbf{B}_k = \nabla^2 f(x_k)$ , we recover the Newton method.